

For the use only of a Registered Veterinary Practitioner or a Hospital or a Laboratory or a Farm.

Flumethrin 1% w/v Pour-On Solution

FlumeticolTM 1% Pour On

Ready-to-use Potent Ectoparasiticide

COMPOSITION:

Each ml contains:
Flumethrin.....1% w/v
Excipients.....q.s.

INDICATIONS:

For the control of ticks (*Ixodes ricinus*), biting lice (*Damalinia bovis*) and sucking lice (*Linognathus vituli*, *Haemotopinus eurysternus*) and psoroptic mange in Cattle.

DOSAGE AND ADMINISTRATION:

Dosage:

Control of ticks and biting lice : 1 ml / 10 kg bodyweight

Control of Sucking lice and mange : 2 ml / 10 kg bodyweight

Treat at 14 day intervals according to tick pressure.

In cases of clinically severe mange a repeat treatment is necessary after 14 days.

Determine the bodyweight accurately before treatment to ensure correct dosage.

ADMINISTRATION:

For topical use.

DIRECTIONS FOR USE:

500 ml : Squeeze the required dose volume into the measuring cup and apply the contents along the back line of the animal from the front of the shoulder to the tail setting.

50 ml /100 ml : Remove the existing cap and seal and use twist-on cap for application. Turn the cap nozzle to open the product and then apply the contents evenly along the back line of the animal from the front of the shoulder to the tail setting.

TARGET SPECIES: Cattle

PHARMACODYNAMIC PROPERTIES:

Pharmacotherapeutic group: Ectoparasiticides for topical use, pyrethroids.

ATCvet code: QP53AC05

PHARMACODYNAMICS:

Flumethrin is an ectoparasiticide of the synthetic pyrethroid group. The synthetic pyrethroids interfere with the sodium channel of nerve cell membranes, resulting in a delay in repolarization of the nerve. Alpha-cyano pyrethroids (type-II pyrethroids) like Flumethrin appear to be much more potent in this regard causing long-lasting trains of repetitive firing in nerve cells. No anti-cholinesterase activity is noted with synthetic pyrethroids.

WARNING & PRECAUTIONS:

- Wear gloves during application.
- Avoid contact with mucous membrane and skin. If the product is in contact, rinse immediately with plenty of water.
- Do not apply to areas of the backline covered with mud or manure.
- Do not smoke, drink or eat while handling the product.
- Keep treated animals away from flame until animal gets dry.
- Wash the hands thoroughly after use.
- It is to be applied to the skin surface only. Do not administer orally or parenterally.

CONTRAINDICATIONS:

Do not administer to animals other than those for which it is indicated.
Hypersensitivity to any of the Ingredients.

ADVERSE EFFECTS:

Slightly irritant to the skin and mucous membranes. If swallowed, sign of intoxication are paresthesia tremor of hyperkinesia. No specific antidote known.

WITHDRAWAL PERIOD:

Meat: 5 days after the latest treatment.

Milk: 10 days after the latest treatment.

TOXICITY:

If swallowed, contact your veterinarian immediately for symptomatic treatment. Only for use by or under the observation of a registered Veterinarian.

STORAGE: Store in a cool and dry place. Protect from light.

PRESENTATION:

- 50 ml / 100 ml HDPE Bottle packed in a Primary Carton along with the Pack Insert .
- 500 ml / 1 Litre HDPE container.

**VETERINARY.
NOT FOR HUMAN USE.
FOR ANIMAL TREATMENT ONLY.
FOR EXTERNAL APPLICATION ONLY.**

SHAKE WELL BEFORE USE.

KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.

Carefully read the accompanying instructions before use.

Veko®

Manufactured & Marketed by:

Veko Care Pvt. Ltd.

Plot No. E-48 & 49, MIDC, Ranjangaon,

Dist-Pune, Maharashtra, India.

Pin Code:412220

Customer care : care@vekocare.com

TM : Trade Mark of Veko Care



VPP1E033