

For the use only of a Registered Veterinary Practitioner or a Hospital or a Laboratory or a Farm.

Diminazene Aceturate, Phenazone Granules with Vitamin B<sub>6</sub> and Vitamin B<sub>12</sub>

## Trypano™ Plus

Granules 2.37 g / 23.7 g Powder for Parenteral Administration.

### COMPOSITION:

Each 2.37 gram sachet contains:

Diminazene Aceturate.....1.05 g  
Phenazone BP.....1.31 g  
Vitamin B<sub>6</sub> (Pyridoxine Hydrochloride) BP.....8 mg  
Vitamin B<sub>12</sub> (Cyanocobalamin) BP.....2 mg  
Excipients.....q.s.

Each 23.7 gram sachet contains:

Diminazene Aceturate.....10.5 g  
Phenazone BP.....13.1 g  
Vitamin B<sub>6</sub> (Pyridoxine Hydrochloride) BP.....80 mg  
Vitamin B<sub>12</sub> (Cyanocobalamin) BP.....20 mg  
Excipients.....q.s.

### INDICATIONS:

Treatment of babesiosis, piroplasmosis and trypanosomiasis in camels, cattle, cats, dogs, goats, horses, sheep and swine.

### ADMINISTRATION:

For Deep Intramuscular Administration.

### DOSEAGE:

1 ml / 20 kg body weight  
(2.37 gram / 300 kg body weight) / (23.7 gram / 3000 kg body weight).  
Dissolve 2.37 gram powder in 15 ml sterile water before use to prepare an injectable solution.  
Dissolve 23.7 gram powder in 150 ml sterile water before use to prepare an injectable solution.  
Use only freshly prepared solutions for injections.

### TARGET SPECIES:

Camels, Cattle, Cats, Dogs, Goats, Horses, Sheep and Swine.

### PHARMACOLOGICAL ACTION:

Diminazene aceturate belongs to the group of aromatic diamidines, with activity against babesiosis, piroplasmosis and trypanosomiasis. Vitamin B<sub>6</sub> and B<sub>12</sub> aid to combat the anaemic effects of the disease through stimulation of red blood cell formation. Vitamin B<sub>6</sub> and B<sub>12</sub> aid to combat the anaemic effects of the disease through stimulation of red blood cell formation.

### WARNING & PRECAUTION:

Do not exceed the recommended dosage.

### CONTRAINDICATIONS:

Do not administer to animals other than those for which it is indicated.  
Hypersensitivity to any of the Ingredients.  
Administration to animals with an impaired renal and / or hepatic function.

### ADVERSE EFFECTS:

Hypersensitivity reactions.  
Salivation, sweating, tremors can occur.

### OVERDOSAGE:

Multiple therapeutic doses can produce severe nervous signs and prominent haemorrhagic and malacic lesions of the cerebellum, midbrain, and thalamus in dogs / cattle.  
After multiple therapeutic doses degenerative fatty changes can occur in liver, kidneys, myocardium and muscles.

### WITHDRAWAL PERIOD:

For Meat: 21 days, For Milk: 4 days.

### STORAGE:

Store below 30°C in a dry place, protected from light.

### PRESENTATION:

10 Sachets of 2.37 g / 23.7g are packed in a primary carton along with the Pack Insert.

**VETERINARY.  
NOT FOR HUMAN USE.  
FOR ANIMAL TREATMENT ONLY.**

**KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.**  
Carefully read the accompanying instructions before use.

Manufactured & Marketed by:

**Veko Care Pvt. Ltd.**  
Plot No. E - 48 & 49, MIDC, Ranjangaon, District - Pune,  
Maharashtra, INDIA. Pin Code : 412220.  
Customer care : care@vekocare.com  
TM : Trade Mark of Veko Care

**Veko** VPP/E/05

