

For the use only of a Registered Veterinary Practitioner or a Hospital or a Laboratory or a Farm.

Quinapyramine Sulphate and Quinapyramine Chloride Powder

QUINACURE™ PLUS

Powder for parenteral administration

COMPOSITION:

Each sachet contains
Quinapyramine Sulphate B.Vet. C.....1.5g
Quinapyramine Chloride B. Vet C....1.0g

INDICATIONS:

QUINACURE PLUS is indicated for prevention & treatment of trypanosomiasis in Camel, Cattle, Horse, Sheep & Goat

DOSE AND ADMINISTRATION:

Dissolve the sachet's content in 15 ml of Water for Injection before use.
1ml/40 kg body weight to be injected subcutaneously.
Use only freshly prepared solutions for injections.
Shake well before use.

Administration: For subcutaneous administration.

WARNING & PRECAUTION:

Determine the body weight accurately before treatment to ensure correct dosage.
Do not exceed the recommended dosage.
Do not inject more than 15 ml by site of injection. In case of higher volume, the dose may be divided into 2 doses with 12 hours.

CONTRAINDICATIONS:

Hypersensitivity to quinapyramine.
Administration to animals with an impaired renal and / or hepatic function.

TARGET SPECIES: Camel, Cattle, Horse, Sheep & Goat

WITHDRAWAL PERIOD: For Meat : 21 days.; For Milk : 4 days.

STORAGE: Store in a dry place below 30°C.

PRESENTATION: 12 Sachets of 2.5 g packed in a primary carton along with the Pack Insert.

**VETERINARY.
NOT FOR HUMAN USE.
FOR ANIMAL TREATMENT ONLY.**

KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.
Carefully read the accompanying instructions before use.

Veko

Manufactured & Marketed By:
VEKO CARE PVT.LTD.

Plot No. E-48 & 49, MIDC, Ranjangaon, District-Pune, Maharashtra,
INDIA. Pin Code:412220.

Customer care:care@vekocare.com TM:Trade Mark of Veko Care

