

For the use only of a Registered Veterinary Practitioner or a Hospital or a Laboratory or a Farm.

## Isometamidium Chloride Hydrochloride Powder

# VEKODIUM™

Granules for Injectable Preparation  
Prevention and Treatment of *Trypanosomiasis*.

#### COMPOSITION:

Each sachet contains:

Isometamidium Chloride Hydrochloride .....1 g  
Excipient (Antipyrine) s.q.f.....6.2 g

#### COMPOSITION:

Each sachet contains:

Isometamidium Chloride Hydrochloride .....125 mg  
Excipient (Antipyrine) s.q.f.....775 mg

#### DESCRIPTION:

Isometamidium is a phenanthridine aromatic amidine with a narrow therapeutic index as both a prophylactic and a therapeutic trypanocidal agent. Isometamidium chloride is used curatively at lower dosage rates, and prophylactically at higher dosage rates.

#### INDICATIONS:

VEKODIUM is indicated for treatment and prevention of trypanosomiasis caused by *Trypanosoma spp.* in Cattle (*incl. zebu*), buffaloes, sheep, goats, camels, horses, donkeys and dogs.

#### DOSAGE AND ADMINISTRATION (6.2g):

- Inject by deep intramuscular route.
- The general dose is 1 ml for 20 to 80 Kg of body weight. (i.e 0.25 to 1mg of Isometamidium/Kg of body weight.)
- Dissolve the content of one sachet of VEKODIUM in 50 ml (2% solution) or 100 ml (1% solution) sterile water for injection.

**DOSAGE AND ADMINISTRATION (775mg):** Dissolve the content of one sachet of VEKODIUM in 6.25 ml (2% solution) or 12.5 ml (1% solution) sterile water for injection.

Species	2% Solution	
	Curative Dose	Preventive Dose
Cattle	2.5 ml/100 Kg	5 ml/100 Kg
Buffaloes	5 ml/100 Kg	5 ml/100 Kg
Sheep, Goats	0.25 ml to 0.5 ml /20 Kg	0.5 ml/20 Kg
Camels	2.5 ml/100 Kg	3.5 ml/100 Kg
Horses, Donkeys	2.5 ml/100 Kg	2.5 ml/100 Kg
Dogs	0.5 ml/10 Kg	0.50 ml/10 Kg

#### WARNING & PRECAUTION:

- Do not inject subcutaneously.
- Do not inject more than 15 ml by site of injection: in case of higher volume, divide the total dose in 2 injections one far from the other.
- Do not inject VEKODIUM with another trypanocide at the same time.

#### CONTRAINDICATIONS:

VEKODIUM should not be administered subcutaneously. Avoid concurrent administration of other trypanocidal drugs. Avoid under dosing.

**SIDE EFFECTS:** Individual sensitivity to VEKODIUM is possible.

**WITHDRAWAL PERIOD:** Meat & Offal : 30 days, Milk : Nil

**STORAGE:** Store below 30°C in a dry place. Protect from light.

#### PRESENTATION:

- 10 Sachets of 6.2 g packed in a primary carton along with the Pack Insert.
- 10 Sachets of 775 mg packed in a primary carton along with the Pack Insert.

**FOR VETERINARY USE. NOT FOR HUMAN USE. FOR ANIMAL TREATMENT ONLY.**

**KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.**

Carefully read the accompanying instructions before use.

Manufactured & Marketed By:

VEKO CARE PVT.LTD.  
Plot No. E-48 & 49, MIDC, Ranjangaon, District-Pune,  
Maharashtra, INDIA. Pin Code:412220.  
Customer care:care@vekocare.com  
TM:Trade Mark of Veko Care

**Veko**

