



A healthy buddy. A happy buddy.

Pack Insert Text

Oxyclozanide and Levamisole Hydrochloride Suspension ***VEmisole – OZ***

For the control of Fluke & Worm by Drench.

COMPOSITION:

Each ml contains :

Oxyclozanide BP 60 mg

Levamisole Hydrochloride BP 30 mg

Approved colours.

Excipients q.s.

INDICATIONS:

For the treatment and control of both gastrointestinal and pulmonary nematode infections, and adult liver fluke infections. The product removes most mature *Fasciolaspp* (flukes) present in the bile ducts of the liver.

DOSAGE AND ADMINISTRATION:

Cattle & Calves : 2.5 ml / 10 kg bodyweight

Sheep & Goats : 0.5 ml / 2 kg bodyweight

Or as directed by the Veterinarian for appropriate dosing programmes.

For Oral Administration.

TARGET SPECIES:

Cattle , Calves , Sheep & Goats .

PHARMACODYNAMIC PROPERTIES:

Pharmacotherapeutic group: Anthelmintic

ATC Vet Code: QP 52 AE 51

PHARMACODYNAMICS:

Levamisole : Is an imidazothiazole that acts by interfering with parasite nerve transmission causing muscular paralysis. It is effective against adult and immature gastrointestinal roundworm and lungworm infections.

Oxyclozanide : Is a salicylanilide, which is mainly active against adult liver flukes. It is distributed to the liver, kidney and intestines and is excreted in the bile.

WARNINGS:

- Determine the bodyweight accurately before treatment to ensure correct dosage.
- Shake the container well before use.
- Keep away from food, drink and animal feeding stuff.
- Do not smoke , drink or eat while handling the product.
- Do not mix with other products.
- Wash the hands thoroughly after use.
- Avoid contact with the drug in the skin and eyes.
- Veterinary advice should be sought if the product does not achieve the desired clinical effect since other diseases, nutritional disturbances or anthelmintic resistance may be involved.
- After treatment, animals should be moved to clean pasture in order to prevent re-infection.



A healthy buddy. A happy buddy.

CONTRAINDICATIONS:

Do not administer to animals other than those for which it is indicated.

Hypersensitivity to any of the Ingredients.

Animals must not be treated within a period of 14 days before or after treatment with organophosphorus compounds.

ADVERSE EFFECTS:

At normal oxyclozanide dose levels, cattle may show slight softening of the faeces with the occasional animal showing increased frequency of defecation and transient inappetence. Rarely, sheep may show an anaphylactic reaction with swelling of the head.

PREGNANCY AND LACTATION:

The product can be safely administered to pregnant or lactating animals. Care should be taken not to exceed the recommended dose.

OVERDOSAGE :

Do not exceed the recommended dosage.

WITHDRAWAL PERIOD:

Meat : 5 days after the latest treatment

Do not use in Cattle/Sheep producing milk intended for human consumption

STORAGE:

Store below 30°C. Protect from light.

PRESENTATION:

1000ml Bottle.

**VETERINARY.
FOR ANIMAL TREATMENT ONLY.
NOT FOR HUMAN USE.**

KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.

Carefully read the accompanying instructions before use.

Manufactured For :

VEKO CARE PVT. LTD.

Room No. 1, Ground Floor , Plot No. E - 47,

M.I.D.C., Ranjangaon , Shirur ,

Tal. Shirur (Pune – Zone 4) - 412220.India

Customer care : care@vekocare.com

TM : Trade Mark of Vekocare