

90.00 mm

For the use only of a Registered Veterinary Practitioner or a Hospital or a Laboratory or a Farm.

Fenbendazole Veterinary Oral Paste BP 10% w/w

Fenland™ 10% Oral Paste

Vanilla
Flavour

A Broad Spectrum Anthelmintic for Dog, Cats, Puppies & Kittens

COMPOSITION:

Each ml contains:
Fenbendazole BP (Vet).....100 mg
Excipients.....q.s.

CATEGORY: A Broad Spectrum Anthelmintic for Dog, Cats, Puppies & Kittens.

INDICATIONS:

Fenland Paste 10 % is indicated for the for the treatment of dogs and cats infected with immature and mature stages of nematodes of the gastro-intestinal and respiratory tracts. The product also has an ovicidal effect on nematode eggs.

Adult dogs and cats:

For the treatment of adult dogs and cats infected with gastrointestinal nematodes & cestodes:

- *Ascarid* spp (*Toxocara canis*, *Toxocara cati* and *Toxascaris leonina*)
- *Ancylostoma* spp • *Trichuris* spp
- *Uncinaria* spp • *Taenia* spp

Puppies and kittens:

For the treatment of puppies and kittens infected with gastrointestinal nematodes and puppies infected with protozoa (*Giardia* spp).

Also for the treatment of dogs infected with lungworm *Ostertus (Filaroides) osleri* or protozoa *Giardia* spp and cats infected with lungworm *Aelurostrongylus abstrusus*.

DOSAGE AND ADMINISTRATION:

The recommended Oral dose to be used is 50 mg fenbendazole per kg bodyweight. 5 ml of **Fenland Paste 10 %** is sufficient to dose up to 10 kg bodyweight as a single dose.

The dosing is as follows:

Body Weight Range of Pets	Dosage	Dose of Fenbendazole
1.1 to 2 Kgs	1.0 ml	100 mg
2.1 to 10 Kgs	5.0 ml	500 mg
10.1 to 20 Kgs	10.0 ml	1.0 g
20.1 to 30 Kgs	15.0 ml	1.5 g
30.1 to 40 Kgs	20.0 ml	2.0 g
40.1 to 50 Kgs	25.0 ml	2.5 g
50.1 to 60 Kgs	30.0 ml	3.0 g

Note : Position locking ring over appropriate mark on plunger.

For the treatment of clinical worm infestations in adult dogs and cats or *Giardia* spp. infections in dogs and puppies, orally administer 50 mg fenbendazole/kg bodyweight daily for 3 days.

For the control of lungworm *Ostertus (Filaroides) osleri* in dogs administer 50 mg fenbendazole/kg bodyweight daily for 7 days. A repeat course of treatment may be required in some cases.

For the control of lungworm *Aelurostrongylus abstrusus* in cats administer 50 mg fenbendazole/kg bodyweight daily for 3 days.

120.00 mm

TARGET SPECIES:

Dog, Cats, Puppies & Kittens.

DIRECTIONS FOR USE:

For oral use.

Before administration, adjust the syringe to the calculated dosage by setting the ring on the plunger. The paste is administered orally by inserting the nozzle of the syringe through the interdental space and depositing the required amount of the paste on the back of the tongue. The animal's mouth should be free of any food. Immediately after administration, elevate the head of the pet for a few seconds to ensure the dose is swallowed.

PHARMACOLOGICAL PROPERTIES:

Pharmacotherapeutic group: Anthelmintics, benzimidazoles and related Substances.

ATCvet code: QP52AC13

PHARMACODYNAMICS:

Fenbendazole is an anthelmintic belonging to the benzimidazole carbamates group. It acts by interfering in the energy metabolism of the nematode. The anthelmintic efficacy is based on inhibition of the polymerisation of tubulin to microtubuli. The anthelmintic affects both adult and immature stages of gastrointestinal and respiratory nematodes. Fenbendazole is metabolised to its sulphoxide, then to sulphone and amines.

PHARMACOKINETICS:

Fenbendazole is only partly absorbed after oral administration and is then metabolised in the liver. Fenbendazole and its metabolites are distributed throughout the body and high concentrations can be found in the liver. The elimination of fenbendazole and its metabolites occurs primarily via the faeces (>90%) and to a smaller extent in the urine and milk.

WARNING & PRECAUTIONS:

- Wear gloves during application.
- Wash the hands thoroughly after use.
- Assess bodyweight as accurately as possible before calculating the dosage.
- Intensive use or misuse of anthelmintics can give rise to resistance. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

OVERDOSAGE:

Benzimidazoles have a high margin of safety and is well tolerated in pets.

CONTRAINDICATIONS: None.

WITHDRAWAL PERIOD: Not Applicable.

STORAGE:

Store below 30°C in a cool and dry place.

Discard after 6 months from the date of the first use.

PRESENTATION:

- 1) 15 ml syringe packed in a Primary Carton along with the Pack Insert.
- 2) 30 ml syringe packed in a Primary Carton along with the Pack Insert.

**FOR VETERINARY USE ONLY.
NOT FOR HUMAN USE.
FOR ANIMAL TREATMENT ONLY.**

KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.

Carefully read the accompanying instructions before use.

Veko®

Manufactured & Marketed by:
VEKO CARE PVT. LTD.

Plot No. E-48 & 49, MIDC, Ranjangaon, Dist-Pune,
Maharashtra, INDIA, Pin Code : 412220.

Customer care : care@vekocare.com

TM : Trade Mark of Veko Care

