

115 mm

120 mm



DoxyKO[®]-20

Doxycycline Hyclate Soluble Powder 20% W/W

**POWDER FOR
ORAL ADMINISTRATION**



Indicated for the prevention and treatment of bacterial enteritis in **Calves, Sheep, Goats, Swine & Poultry**

Sachet of **100g**

DoxyKO[®]-20

Doxycycline Hyclate Soluble Powder 20% W/W

COMPOSITION:

Each gram contains:
Doxycycline Hyclate BP....200 mg
Excipients.....qs

DESCRIPTION:

Doxycycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria, like *Bordetella*, *Campylobacter*, *E. coli*, *Haemophilus*, *Pasteurella*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. Doxycycline is also active against *Chlamydia*, *Mycoplasma* and *Rickettsia* spp. The action of doxycycline is based on inhibition of bacterial protein synthesis. Doxycycline has a great affinity to the lungs and is therefore especially useful for treatment of bacterial respiratory infections.

INDICATIONS:

DoxyKO-20 is indicated for the prevention & treatment of Gastrointestinal and respiratory infections caused by doxycycline sensitive micro-organisms, like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. in Calves, Sheep, Goats, Swine and Poultry.

CONTRAINDICATIONS:

Hypersensitivity to tetracyclines.
Administration to animals with a seriously impaired hepatic function.
Concurrent administration of penicillins, cephalosporins, quinolones and cycloserine.
Administration to animals with an active microbial digestion.

SIDE EFFECTS

Discoloration of teeth in young animals may occur.
Hypersensitivity reactions may occur.

DOSAGE AND ADMINISTRATION:

Powder For oral administration:

Calves, Goats and Sheep:

Twice daily 5 g per 200 kg body weight for 3 - 5 days.

Poultry and Swine:

100 gram per 200 - 400 litres of drinking water for 3 - 5 days.

Note: for pre-ruminant calves, lambs and kids only.

WITHDRAWAL PERIOD:

Meat :

Calves, Goats and Sheep : 14 days after the latest treatment.

Swine : 8 days after the latest treatment.

Poultry : 7 days after the latest treatment.

Milk : 0 days ; Not for use in animals from which milk or eggs are produced for human consumption.

STORAGE:

Store below 30°C in a dry place.

Mfg. Lic. No. : MH/102594

Batch No. :

Unvarnish Area
26 x 12 mm

Mfg. Date :

Exp. Date :



8 19 06 1021 18 05 15

Manufactured & Marketed By:
VEKO CARE PVT. LTD.

Plot No. E - 48 & 49, MIDC,
Ranjangaon, District - Pune,
Maharashtra, INDIA. Pin Code : 412220.
Customer care : care@yekocare.com
® : Registered Trade Mark of Veko Care

VPL18037