

115 mm



DoxyKO-50[®]

Doxycycline Hyclate Soluble Powder 50% W/W

**POWDER FOR
ORAL ADMINISTRATION**



Indicated for the prevention and treatment of bacterial enteritis in **Calves, Sheep, Goats, Swine and Poultry**

**FOR VETERINARY USE ONLY.
NOT FOR HUMAN USE.
FOR ANIMAL TREATMENT ONLY.**

Sachet of
100g

120 mm

DoxyKO-50[®]

Doxycycline Hyclate Soluble Powder 50% W/W

COMPOSITION:

Each gram contains:
Doxycycline Hyclate BP.....50% w/w
Excipients.....qs

DESCRIPTION:

Doxycycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria, like *Bordetella*, *Campylobacter*, *E. coli*, *Haemophilus*, *Pasteurella*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. Doxycycline is also active against *Chlamydia*, *Mycoplasma* and *Rickettsia* spp. The action of doxycycline is based on inhibition of bacterial protein synthesis. Doxycycline has a great affinity to the lungs and is therefore especially useful for treatment of bacterial respiratory infections.

INDICATIONS:

DoxyKO-50 is indicated for the prevention & treatment of Gastrointestinal and respiratory infections caused by doxycycline sensitive micro-organisms, like *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. in Calves, Sheep, Goats, Swine and Poultry.

CONTRAINDICATIONS:

Hypersensitivity to tetracyclines.
Administration to animals with a seriously impaired hepatic function.
Concurrent administration of penicillins, cephalosporins, quinolones and cycloserine.
Administration to animals with an active microbial digestion.

SIDE EFFECTS

Discoloration of teeth in young animals may occur.
Hypersensitivity reactions may occur.

DOSAGE AND ADMINISTRATION:

Powder For oral administration:

Calves, Goats and Sheep:

Twice daily 1 g per 100 kg body weight for 3 - 5 days.

Poultry and Swine:

100 gram per 500 - 1000 litres of drinking water for 3 - 5 days.

Note: for pre-ruminant calves, lambs and kids only.

WITHDRAWAL PERIOD:

Meat :

Calves, Goats and Sheep : 14 days after the latest treatment.

Swine : 8 days after the latest treatment.

Poultry : 7 days after the latest treatment.

Milk : 0 days; Not for use in animals from which milk or eggs are produced for human consumption.

STORAGE:

Store below 30°C in a dry place.

**KEEP OUT OF REACH OF CHILDREN & PETS,
AWAY FROM FOOD.**

Carefully read the accompanying instructions before use.

Mfg. Lic. No. : MH/102594

Batch No. : Unvarnish Area
Mfg. Date : 26 x 10 mm
Exp. Date :



8 906102 160522

Manufactured & Marketed By:
VEKO CARE PVT. LTD.
Plot No. E - 48 & 49, MIDC,
Ranjangaon, District - Pune,
Maharashtra, INDIA. Pin Code : 412220.
Customer care : care@vekocare.com
® : Registered Trade Mark of Veko Care

