

115 mm

120 mm




OtcRxTM-50

Oxytetracycline Hydrochloride Soluble Powder 50% W/W

100 g

POWDER FOR ORAL ADMINISTRATION



Gastrointestinal and respiratory infections caused by oxytetracycline sensitive bacteria in
Calves, Sheep, Goats, Swine and Poultry

115 mm

120 mm

OtcRxTM-50

COMPOSITION:

Each gram contains:
Oxytetracycline Hydrochloride BP.....500 mg

DESCRIPTION:

Oxytetracycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like Bordetella, Bacillus, Corynebacterium, Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella, Staphylococcus and Streptococcus spp. and Mycoplasma, Rickettsia and Chlamydia spp. The mode of action of oxytetracycline is based on inhibition of bacterial protein synthesis. Oxytetracycline is mainly excreted in urine and to a lesser degree in bile and in lactating animals in milk.

INDICATIONS:

OtcRx- 50 is indicated for the prevention & treatment of Gastrointestinal and respiratory infections caused by oxytetracycline sensitive bacteria like Bordetella, Bacillus, Corynebacterium, Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella, Staphylococcus and Streptococcus spp. and Mycoplasma, Rickettsia and Chlamydia spp. in calves, goats, poultry, sheep and swine.

DOSAGE AND ADMINISTRATION:

Powder For oral administration:
Calves, Goats and Sheep:
Twice daily 1 g per 50-100 kg body weight for 3 - 5 days.
Poultry and Swine:
100 gram per 500 litres of drinking water for 3 - 5 days.
Note: for pre-ruminant calves, lambs and kids only.

WARNING & PRECAUTION:

- Do not exceed the recommended dosage.
- Determine the bodyweight accurately before treatment to ensure correct dosage.
- Wash the hands thoroughly after use.
- In case of accidental ingestion by any Human being, seek medical advice immediately and show the package leaflet or the label to the physician.

TARGET SPECIES: Calves , Sheep, Goats, Swine and Poultry

Mfg. Lic. No. : MH/102594

Batch No. :

Mfg. Date :

Exp. Date :

Unvannish Area
20 X 9 mm

CONTRAINDICATIONS:

Hypersensitivity to tetracyclines.
Administration to animals with a seriously impaired hepatic function.
Concurrent administration of penicillins, cephalosporins, quinolones and cycloserine.
Administration to animals with an active microbial digestion.

WITHDRAWAL PERIOD:

Meat :
Calves, Goats , Sheep and Swine : 8 days after the latest treatment.
Poultry : 6 days after the latest treatment.
Eggs : 6 days after the latest treatment.

STORAGE: Store below 30 °C in a dry place.

**FOR VETERINARY USE ONLY.
NOT FOR HUMAN USE.
FOR ANIMAL TREATMENT ONLY.**

KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.
Carefully read the accompanying instructions before use.



8 19 06 1021 180531



MANUFACTURED & MARKETING BY:

VEKO CARE PVT. LTD.
Plot No. E - 48 & 49, MIDC,
Ranjangaon , District - Pune ,
Maharashtra , INDIA. Pin Code : 412220.
Customer care :care@vekocare.com
TM : Trade Mark of Veko Care

