

For the use only of a Registered Veterinary Practitioner or a Hospital or a Laboratory or a Farm.

Oxyclozanide Tablet 750 mg

CLOZE™ 750

A Broad Spectrum Anthelmintic for Oral use.

COMPOSITION:

Each uncoated tablet contains:-
Oxyclozanide BP (Vet).....750 mg
Excipients.....q.s.

INDICATIONS:

CLOZE 750 is indicated in the treatment of acute and chronic fasciolosis (Fasciola hepatica and F. gigantica) in Cattle, Buffaloes, Sheep and Goats caused by both mature and immature flukes. It is also recommended in the treatment of amphistomiasis in bovines and for tapeworms (Moniezia spp.) in Sheep.

DOSAGE AND ADMINISTRATION:

- Sheep /Goat : 1 bolus per 50 kg body weight.
- Cattle/ Buffaloes : 1 bolus per 75 kg body weight .
Or as directed by the Veterinarian.

Administration:

To be administered orally by means of a balling gun, or to be dissolved in water and given orally.

TARGET SPECIES:

Sheep/Goat /Cattle/ Buffaloes.

PHARMACODYNAMIC PROPERTIES:

Pharmacotherapeutic group: Anthelmintics, Phenol derivatives, incl. salicylanilides, oxyclozanide
ATC vet code: QP52AG06

PHARMACODYNAMICS:

Oxyclozanide is an anthelmintic of the salicylanilide group. The mechanism of action of oxyclozanide is the uncoupling of oxidative phosphorylation in flukes.
Oxyclozanide has flukicidal activity against the adult stage of Fasciola hepatica. Its efficacy against cestodes is limited to the removal of segments of the tapeworm Moniezia.

WARNING & PRECAUTIONS :

- Do not exceed the recommended dosage.
- Determine the bodyweight accurately before treatment to ensure correct dosage.
- Wash the hands thoroughly after use.

CONTRAINDICATIONS:

Do not administer to animals other than those for which it is indicated.
Hypersensitivity to any of the Ingredients.

PREGNANCY AND LACTATION:

Care should be taken not to exceed the recommended dose during pregnancy.

DRUG INTERACTIONS:

None Known.

ADVERSE EFFECTS :

At normal oxyclozanide dose levels, cattle may show slight softening of the faeces with the occasional animal showing increased frequency of defecation and transient inappetence.

OVERDOSAGE:

The effects of oxyclozanide over dosage are possibly diarrhoea, inappetence and loss of weight in cattle.

WITHDRAWAL PERIOD:

- Meat : 14 days after the latest treatment.
- Milk : 24 hours after the latest treatment.

STORAGE:

Store in a cool & dry place below 25°C .

**FOR VETERINARY USE.
FOR ANIMAL TREATMENT ONLY
NOT FOR HUMAN USE**

KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.

Carefully read the accompanying instructions before use.

Veko®

**Manufactured & Marketed By:
VEKO CARE PVT. LTD.**

Plot No. E - 48 & 49, MIDC ,
Ranjangaon , District - Pune , Maharashtra ,
INDIA, Pin Code : 412220.
Customer care : care@vekocare.com
TM : Trade Mark of Veko Care.

XXX