

90.00 mm

120.00 mm

For the use only of a Registered Veterinary Practitioner or a Hospital or a Laboratory or a Farm.

## Sucralfate Oral Gel

# STOPCER™ ORAL GEL

### COMPOSITION:

Each ml contains:

Sucralfate.....240 mg

Excipients.....q.s.

*Vanilla Flavoured-For Pets*

*Green Apple Flavoured-For Horses*

### INDICATIONS:

**STOPCER ORAL GEL** is an antiulcer drug that is indicated for the treatment and prevention of gastric and duodenal ulcers, reflux esophagitis and gastropathies caused by drug administration.

### DOSAGE AND ADMINISTRATION:

**Canines:** 500 mg in small dogs and 1 g in large dogs.

**Horses:** 10 to 20 mg/kg of body weight. Equivalent to 1 ml for every 12 to 24 kg. every to 12 hrs.

The duration of the treatment will be according to the criteria of the Veterinarian.

For oral use only.

### DIRECTIONS FOR USE:

- It is intended for use only with the dispensing syringe designed for this product.
- Set the correct dose rate by turning the plunger lock.
- Remove the cap from the end of the syringe.
- Canines, Horses's Mouth must be free of food.
- Insert the syringe nozzle into the side of the mouth, in the gap between the front and back teeth. Advance the syringe as far as possible into the mouth, and deposit the paste onto the back of the tongue by pressing the plunger completely to give the measured dose.
- Immediately after administration, raise the Vet's head & hold for a few seconds to ensure the full dose is swallowed.
- Replace the syringe cap

**ADMINISTRATION:** Oral

**TARGET SPECIES:** Cats, Dogs & Horses .

### PHARMACODYNAMIC PROPERTIES :

**Pharmacotherapeutic group:** Alimentary tract and metabolism, ATC code: A02B X02

### PHARMACODYNAMICS :

Sucralfate is a sucrose-sulfate-aluminium complex that acts as a local protectant of the digestive mucosa, binds to the surface of ulcers and also acts as a buffer due to its cytoprotective properties.

### PHARMACOKINETICS :

Sucralfate is only minimally absorbed from the gastro-intestinal tract. The small amounts that are absorbed are excreted primarily in the urine. Absorption of aluminium from sucralfate may be increased in patients on dialysis or with renal dysfunction .

### WARNING & PRECAUTION :

#### Special precautions for use in animals:

- Do not use the same syringe in more than one animal.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- Wear gloves during application.
- Wash the hands thoroughly after use.
- Do not eat, drink or smoke while handling this product.
- Assess bodyweight as accurately as possible before calculating the dosage.
- In case of accidental eye contact, rinse immediately with plenty of water.
- In case of accidental ingestion or eye irritation, seek medical advice and show the package leaflet or label to the physician.

### CONTRAINDICATIONS:

Hypersensitivity to any of the Ingredients.  
Do not administer to animals other than those for which it is indicated.

### ADVERSE EFFECTS:

None.

### PREGNANCY AND LACTATION:

It should be used during pregnancy only if clearly needed.  
Lactation: It is not known whether this drug is excreted in milk. Caution should be exercised when it is administered .

### WITHDRAWAL PERIOD:

Not Applicable.

### STORAGE:

Store below 25°C. Discard after 6 months from the date of the first use.

### PRESENTATION:

30ml HDPE Syringe packed in a Primary Carton along with the Pack Insert.

FOR VETERINARY USE.  
FOR ANIMAL TREATMENT ONLY.  
NOT FOR HUMAN USE.

### KEEP OUT OF REACH OF CHILDREN & PETS, AWAY FROM FOOD.

Carefully read the accompanying instructions before use.

Manufactured by:

**VEKO CARE PVT. LTD.**

Plot No. E - 48 & 49, MIDC ,Ranjangaon ,  
District - Pune, Maharashtra, INDIA. Pin Code : 412220.

Customer care : care@vekocare.com

TM : Trade Mark of Veko Care

**Veko®**

